


# Comparative Genomics of Early-Diverging *Brucella* Strains Reveals a Novel Lipopolysaccharide Biosynthesis Pathway

Alice R. Wattam,<sup>a</sup> Thomas J. Inzana,<sup>b</sup> Kelly P. Williams,<sup>c</sup> Shrinivasrao P. Mane,<sup>a</sup> Maulik Shukla,<sup>a</sup> Nalvo F. Almeida,<sup>d</sup> Allan W. Dickerman,<sup>a</sup> Steven Mason,<sup>a</sup> Ignacio Moriyón,<sup>e</sup> David O'Callaghan,<sup>f,g</sup> Adrian M. Whatmore,<sup>h</sup> Bruno W. Sobral,<sup>a</sup> Rebekah V. Tiller,<sup>i</sup> Alex R. Hoffmaster,<sup>i</sup> Michael A. Frace,<sup>j</sup> Cristina De Castro,<sup>j</sup> Antonio Molinaro,<sup>j</sup> Stephen M. Boyle,<sup>b</sup> Barun K. De,<sup>i</sup> and João C. Setubal<sup>a,k</sup>

Virginia Bioinformatics Institute, Virginia Tech, Blacksburg, Virginia, USA<sup>a</sup>; Center for Molecular Medicine and Infectious Diseases, Virginia-Maryland College of Veterinary Medicine, Virginia Tech, Blacksburg, Virginia, USA<sup>b</sup>; Sandia National Laboratories, Livermore, California, USA<sup>c</sup>; Faculdade de Computação, Universidade Federal de Mato Grosso do Sul, Campo Grande, MS, Brazil<sup>d</sup>; Instituto de Salud Tropical y Departamento de Microbiología y Parasitología, Facultad de Medicina, Universidad de Navarra, Edificio de Investigación, Universidad de Navarra, c/Irunlarrea, Pamplona, Spain<sup>e</sup>; INSERM U1047, UFR Médecine, Nîmes, France<sup>f</sup>; Université Montpellier 1, UFR Médecine, Nîmes, France<sup>g</sup>; Department of Bacteriology, Animal Health & Veterinary Laboratories Agency, Woodham Lane, Addlestone, United Kingdom<sup>h</sup>; Centers for Disease Control and Prevention, Atlanta, Georgia, USA<sup>i</sup>; Department of Chemical Sciences, Università di Napoli Federico II, Complesso Universitario Monte Santangelo, Naples, Italy<sup>j</sup>; and Department of Biochemistry, Institute of Chemistry, University of São Paulo, São Paulo, Brazil<sup>k</sup>

**V**olume 3, no. 5, doi:10.1128/mBio.00246-12, 2012. After careful review of Fig. 3, the accuracy of the electrophoretic profiles of the control lipopolysaccharides (LPSs) was called into question compared to profiles of similar LPSs analyzed by M. S. Zygmunt et al. (Clin. Vaccine Immunol. 19:1370–1373, 2012). After further gel experimentation, it was confirmed that the profiles of the control LPSs in Fig. 3 only were not correct. The revised Fig. 3 (below) shows the correct profiles of the LPSs of strains 2308 and BO1. We thank Axel Cloeckert for notifying us of this issue.



**FIG 3** Electrophoretic profiles of the LPSs produced by *B. abortus* 2308, *B. inopinata* strain BO1, and *B. inopinata*-like strain BO2. Lanes: A, *B. abortus* strain 2308 LPS isolated from the phenol phase (80  $\mu$ g); B, strain BO1 LPS isolated from the phenol phase (30  $\mu$ g); C, strain BO2 LPS isolated from the phenol phase (25  $\mu$ g); D, strain BO2 LPS isolated from the aqueous phase (25  $\mu$ g). Strain 2308 and BO1 LPSs were not present in aqueous-phase extracts (data not shown).

**Published** 6 November 2012

**Editor** Paul Keim, Northern Arizona University

**Citation** Wattam AR, Inzana TJ, Williams KP, et al. 2012. Comparative genomics of early-diverging *Brucella* strains reveals a novel lipopolysaccharide biosynthesis pathway. mBio 3(6):e00388-12. doi:10.1128/mBio.00388-12

**Copyright** © 2012 Wattam et al. This is an open-access article distributed under the terms of the Creative Commons Attribution-Noncommercial-Share Alike 3.0 Unported License, which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

Address correspondence to Thomas J. Inzana, tinzana@vt.edu.